



Group 5.1.1

GROUP 5 LEAVES COMPOUND: PINNATE or PALMATE, WITH 2 or MORE LEAFLETS

GROUP 5.1 LEAFLETS 2

GROUP 5.1.1 DOMATIA PRESENT


See also: *Arytera dictyoneura* in Group 5.1.4A,
Elattostachys bidwillii in Group 5.8.1,
Guioa acutifolia in Group 5.8.1,
Toechima tenax in Group 5.8.1 and
Elattostachys xylocarpa in Group 5.8.2A.

Arytera distylis

TWIN-LEAVED COOGERA

Small to medium-sized tree to 24 m high, usually less than 10 m high, in STRf and DRf; north from Woodburn to Maryborough district. **New growth** limp, pink-red. **Leaflets** mostly 2, lanceolate to oblanceolate, 3–10 cm long, apex bluntly pointed or notched, margins entire, hairless, glossy above, net veins obscure above, distinct below; **domatia** several (2–11), small, in vein angles on lower surface as hairless raised cavities. **Fruit** a capsule, ovoid or 2-lobed, 1–1.8 cm long, yellowish; seeds 1 or 2, black, each enveloped in a red fleshy aril. Family Sapindaceae

NSW, Qld



Mischocarpus anodontus

VEINY PEAR-FRUIT

Shrub or small tree to 10 m high, in DRf; north from Lismore. **Leaflets** 2 or 4, broad-elliptic to ovate, 5–11 cm long, apex bluntly pointed or notched, margins entire, hairless, upper surface slightly glossy, net veins prominent especially in dried leaves; **domatia** large, usually 1, 2 or 3 present as hairless pocket-like cavities on lower surface. **Fruit** a capsule, pear-shaped, 3-angled, 2–3 cm long (including basal stalk of 7–10 mm), red, orange or yellow, hairy; seeds 1 or 2, dark brown, aril almost completely covering seeds, purplish or pinkish red. Family Sapindaceae

NSW, Qld




Mischocarpus australis (*Mischocarpus sundaicus*)

RED PEAR-FRUIT

Small to medium-sized tree to 25 m high, in STRf; north from Minmi (Newcastle area) to Mt Morgan (S of Rockhampton). **Branchlets** and petioles brown. **Leaflets** 2 or 4, glossy, thick and tough, elliptic to oblong-elliptic, 5–15 cm long, apex acuminate to a fine point, margins entire and wavy, leathery, hairless, dark green, upper surface ± glossy; secondary veins distinctly raised on lower surface; **domatia** small but inconspicuous. **Fruit** a capsule, ovoid, 1–1.7 cm long (including basal stalk of 3–6 mm), red to orange, hairy inside; seeds 3–6, black, with a purplish fleshy aril. Family Sapindaceae

NSW, Qld




Mischocarpus ailae (previously included in *M. lachnocarpus* s. lat.)

CALDERA PEAR-FRUIT

Tall shrub or small tree to 10 m high, in cool STRf on the Nightcap Range and Springbrook Plateau. **Branchlets**, buds, petioles and the veins on the lower surface of leaflets covered with rusty brown hairs. **Leaflets** usually 2, or rarely 3 or 4, narrow-elliptic to oblong-elliptic, 5–14 cm long, apex notched, usually with a short bristle, margins entire, leathery to stiff; upper surface green and glossy, lower surface paler; **domatia** present as small hairy pockets or flaps, sometimes inconspicuous. **Fruit** a capsule, pear-shaped, 1.3–2 cm long (including basal stalk of 3–5 mm), 2- or 3-lobed, hairy, red or orange; seeds lack, with an orange or yellow fleshy aril, partly covering seed.

Family Sapindaceae

NSW, Qld



Cupaniopsis wadsworthii

DUCKFOOT

Shrub or small tree to 3 m high, often multistemmed; in STRf and DRf; chiefly north from the Miriam Vale district, Qld, to Magnetic Island (Townsville), N Qld. **Leaflets** mostly 2 or 4, or sometimes 6 in juveniles, broadly cuneate, 2–7.5 cm long, apex concavely truncate, apiculate, base tapered, margins usually entire (sometimes broadly toothed at apex in juveniles), surfaces glossy, hairless except for some scattered hairs on veins on lower surface; **domatia** small, in angles of lateral veins as pocket-like cavities. **Fruit** a capsule, obovoid, 1.5–2 cm long, 3-angled, hairy; **seeds** 1–3, black; partly covered by orange aril. Family Sapindaceae

Qld only

